

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Monsanto Company

Discrete, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FLES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN DUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY ECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS A CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF THE 184 STATE 1542, AS AMENDED, 7 U.S.C. 2321 ET SEO.)

WHEAT, COMMON

'Jagalene'

In Tratimony Alectron, I have hereunto set my hand and caused the seal of the Alant Antiety Frotestion Office to be affixed at the City of Washington, D.C. this sixteenth day of September, in the year two thousand two.

Allest:

Jamfahl.

Commissioner

Plant Variety Protection Office Sgricultural Marketing Service Hereman Hericulture

| | | | Local Kepro | duction of FORM - OIMS NO. 0381-0033 |
|---|------------------------------|--|--------------------------|---|
| U.S. DEPARTMENT OF AGRICULTURE | | The following statements are mo | ade in accordance wit | th the privacy Act of |
| AGRICULTURAL MARKETING SER VICE SCIENCE DIVISION - PLANT VARIETY PROTECTION O | FFICE | 1974 (5 U.S.C. 552a) | | |
| | | Application is required in order | to determine if a pla | nt variety protection |
| APPLICATION FOR PLANT VARIETY PROTECTION (Instructions and information collection burden statement on re | | certificate is to be issued (7 U.S. until certificate is issued (7 U.S. | | n is held confidential |
| 1. NAME OF APPLICANT(S) (as it is to appear on the Certificate) | | 2. TEMPORARY DESIGNATION | N OR | 3. VARIETY NAME |
| Monsanto Company | | EXPERIMENTAL NUMBER W98-362 | | Jagalene |
| 4. ADDRESS (Street and No. or R.F.D. No., City, State, and Zip Code) | | 5. TELEPHONE (include area co | oda) | Jagarene |
| 4. ADDRESS (Sireel and No. 01 K.F.D. No., Chy, state, and Esp Code) | | J. TELEFRONE (Include area co | | PATONIMOR A A A A |
| 700 Chesterfield Parkway North | | 636-737-6089 | 4 | °0°2°0016(|
| St. Louis, Missouri 63198 | | | | |
| | | 6. FAX (include area code) | | P DATE |
| | | | | I , , , , , |
| | | 636-737-7250 | | ½ 5/23/zocz |
| 7. GENUS AND SPECIES NAME | 8. FAMILY NAME (Bot | | | G. FILING AND EXAMINATION FEE: |
| | • | omeur) | | FILING AND EXABITRATION FEE. |
| Triticum aestivum | Gramineae | | | 1 2705.00 |
| 9. CROP KIND NAME (common name) | | | | S DATE |
| Hard Red Winter Wheat | | | | 5/23/200Z |
| 10. IF THE APPLICANT NAMED IS NOT A "PERSON", GIVE FORM OF ORGAN | IIZATION (composition | partnership association etc.) (com | mon name | E |
| | ,aariitoi (to.po.ano,, | , | ,. | 1 CERTIFICATION FEE. |
| Corporation | | | | 1 フィル |
| 11. IF INCORPORATED, GIVE STATE OF INCORPORATION | | 12. DATE OF INCORPORATION | 1 | D DATE |
| Delaware | | 1933 | | 7////// |
| 13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO | SERVE IN THIS APPLI | CATION AND RECEIVE ALL PA | PERS | 14. TELEPHONE (include area code) |
| Ms. Sally Metz | OLK III III III III | Dr. Rollin Sears | 1 200 | <u> </u> |
| 700 Chesterfield Parkway North ANI |) | 6515 Ascher Road | | 636-737-6089 |
| St. Louis, Missouri 63198 | | Junction City, Kansas | 66441 | 15. FAX (include area code) |
| | | | | 636-737-7250 |
| | | | | |
| CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (folional folion). X Exhibit A. Origin and Breeding History of the Variety | ow instructions on revers | e) | | |
| b. X Exhibit B. Statement of Distinctness | | | | |
| c. X Exhibit C. Objective Description of the Variety | | | | |
| d. X Exhibit D. Additional Description of the Variety | | | | |
| e. X Exhibit E. Statement of the Basis of the Applicant's Owne | ership | | | |
| f. X Voucher Sample (2,500 viable untreated seeds, or, for tuber prop | agated varieties veification | that tissue culture will be deposited and | maintained in a public r | epository) |
| g. X Filing and Examination Fee (\$2,450), made payable to "T | reasurer of the United S | tates" (Mail to PVPO) | 1-6 | |
| 17. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE SOLD | BY VARIETY NAME | ONLY, AS A CLASS OF CERTIFI | ED SEED? (See Section | 83 (a) of the Plant Variety Protection Act) |
| X YES (if "yes", answer items 18 and 19 below) | | NO (if 'no", go to item | 20) | |
| 18. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE LIMIT GENERATIONS? | TED AS TO NUMBER (| 19. IF 'YES' TO ITEM 18, WHICH | H CLASSES OF PROD | OUCTION BEYOND BREEDERS SEED? |
| YES X | NO | FOUNDATION | REGISTERED | CERTIFIED |
| | | | | |
| 20. HAS THE VARIETY OR A HYBRID PRODUCED FROM THE VARIETY BEEI YES (iF "YES", give names of countries and dates) | N RELEASED, USED, C | PFFERED FOR SALE, OR MARKE NO | ETED IN THE U.S. OF | OTHER COUNTRIES? |
| [120 (il 120) give names of countries and dates) | | 140 | | |
| | | | | |
| The applicant(s) declare that a viable sample of basic seed of the variety will be fur. applicable, or for a tuber propagated variety a tissue culture will be deposited in a p | | | | such regulations as may be |
| The undersigned applicant(s) is(are) the owner(s) of this sexually reproduced or tub | | | | as required in |
| Section 41, and is entitled to protection under the provisions of Section 42 of the Pla | | ., | ,, 5.0010 0 | |
| Applicant(s) is(are) informed that false representation herein can jeopardize protect | on and result in penalties | | | · |
| SIGNATURE OF APPLICANT (Owneds 1) | | SIGNATURE OF APPLICA | ANT (Owner(s)) | |
| NAME (Please print or tipe) | | NAME (Please print or type | e} | |
| Sally Metz | | The state of the s | • | |
| CAPACITY OR TITLE DATE | ×1 /0 0 | CAPACITY OR TITLE | | DATE |
| Director Wheat Technology | pril 29 20 | 0 <u>/</u> L | | |
| SD-470 (O4-95) | 1/ // | 7 | rea for instructions and | l information collection burden statement) |

Exhibit A. Origin and Breeding History of Jagalene

The cross involving Abilene/Jagger was made in 1994 in the greenhouse at Berthoud, CO. and designated 94x1054. F₁ plants were grown in the greenhouse and F₂ seed planted at Nardin, OK in 1996. The F₂ plot was harvested in bulk because of short plant height and better disease resistance, cleaned with a gravity table and the F₃ seed spaced planted at Berthoud during the fall of 1997. During the spring of 1998 individual space plants were selected based upon plant height, stem strength, head size and yield potential. The F₄ row was designated 97PRY1 151-163 and planted in observation trials throughout Kansas, Oklahoma and Nebraska. Based upon improved disease resistance, short plant height and good test weight patterns the line was selected for yield trials and designated W98-362. In 1999 W98-362 was tested in advanced yield trials at 6 locations. In 2000 the line was tested in elite trials at 14 locations and in 2001 W98-362 was tested 18 locations in the southern Great Plains. In 1999 48 head rows were evaluated for uniformity and plant type, 12 rows were harvested and grown as initial breeders seed. From this increase 6 small plots were bulked to form the breeders seed of W98-362 (Jagalene).

Jagalene has been uniform and stable since 2000. Less than 0.8% of the plants were rogued from the Breeders seed increase in 2001. Approximately 87% of the rogued variant plants were taller height wheat plants (8 to 15 cm) and approximately 3% were awnletted wheat plants. Up to 1% variant plants may be encountered in subsequent generations.

Exhibit B. Statement of Distinctness

Jagalene is most similar to the hard red winter wheat 'Abilene'. However, it can be easily distinguished by the following morphological characteristics:

- Jagalene has a medium glume width, (Berthoud, CO 2000 and 2001). Abilene has a narrow shoulder on the glume, (Berthoud, CO 2000 and 2001).
- Jagalene has a long glume length, (Berthoud, CO 2000 and 2001). Abilene has a medium glume length, (Berthoud, CO 2000 and 2001).

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE DIVISION BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY

WHEAT (Triticum Spp.)

| NAME OF APPLICANT(S) | FOR OFFICIAL USE ONLY |
|--|--|
| Monsanto Company | PVPO NUIZEU 0 2 0 0 1 6 0 |
| ADDRESS (Street and No. or R.F.D. No., City, State, and Zip Code) 700 Chesterfield Parkway North St. Louis, Missouri 63198 | NAME OR EXPERIMENTAL DESIGNATION Jagalene |
| Place the appropriate number that describes the varietal character of this variety in the boxes below. Place a zero in the first box when number is either 99 or less or 9 or less respectively. Data for quantitative plant chara minimum of 100 plants. Comparative data should be determined from varieties entered in the same trial. Royal Hortistandard may be used to determine plant colors; designate system used. Please answer all questions for your variety; lack of response may delay progress of your application. | cters should be based on a sultural Society or any recognized |
| 1. KIND: | |
| 1 1=Common 2=Durum 3=Club 4=Other (specify) | |
| 2. VERNALIZATION: | ······································ |
| 2 1=Spring 2=Winter 3=Other (specify) | **** |
| 3. COLEOPTILE ANTHOCYANIN: | |
| 1 1=Absent 2=Present | |
| 4. JUVENILE PLANT GROWTH: | |
| 2 1=Prostrate 2=Semi-erect 3=Erect | |
| 5. PLANT COLOR (boot stage): | |
| 3 1 = Yellow-Green 2 = Green 3 = Blue-Green | |
| 6. FLAG LEAF (boot stage): | |
| $1 = \text{Erect} \qquad 2 = \text{Recurved}$ | |
| 2 1 = Not Twisted 2 = Twisted | |
| 7. EAR EMERGENCE: | |
| 0 0 Number of Days Earlier Than | * |
| 0 3 Number of Days Later Than Jagger | * |
| 3. ANTHER COLOR: | |
| $1 = YELLOW \qquad 2 = PURPLE$ | |
| . PLANT HEIGHT (from soil to top of head, excluding awns): | |
| 0 0 cm Taller Than | * |
| 0 5 cm Shorter Than Jagger | * |
| * Relative to a PVPO-Apprved Commercial Variety Grown in the Same Trial | |

| - | | |
|---------------------------|------------|---|
| 10. | | EM: ANTHOCYANIN |
| 1 | | 1= Absent 2=Present |
| | В. | WAXY BLOOM |
| 2 | | 1=Absent 2=Present |
| | C. | HAIRINESS (last internode of rachis) |
| 2 | | 1=Absent 2=Present |
| | D. | INTERNODE (specify number) |
| 1 | | 1=Hollow 2=Semi-solid 3=Solid |
| | E. | PEDUNCLE |
| 1 | | 1=Erect 2=Recurved |
| 2 | 6 | cm Length |
| | | AD (at Maturity): |
| $\overline{}$ | A. | DENSITY |
| 2 | n | 1=Lax 2=Middense 3= Dense |
| 2 | ь. | SHAPE 1 = Tapering 2= Strap 3 = Clavate 4 = Other (specify) |
| | ٦. | CURVATURE |
| 2 | ∪ , | 1 = Erect 2 = Inclined 3 = Recurved |
| | Э. | AWNEDNESS |
| 4 | | 1 = Awnless 2 = Apically Awnletted 3 = Awnletted 4 = Awned |
| | | JMES (at Maturity): |
| | A . | COLOR |
| 1 | | 1 = White 2 = Tan 3 = Other (specify) |
| | 3. | SHOULDER |
| _2_ | , | 1 = Wanting 2 = Oblique 3 = Rounded 4 = Square 5 = Elevated 6 = Apiculate BEAK |
| 3 | ٠. | 1 = Obtuse 2 = Acute 3 = Acuminate |
| | 1 | LENGTH |
| 3 | ٠. | 1 = Short (ca. 7mm) 2 = Medium (ca. 8mm) 3 = Long (ca. 9mm) |
| 4 | ٠ . ' | WIDTH |
| 2 | | 1 = Narrow (ca. 3mm) 2 = Medium (ca. 3.5mm) 3 = Wide (ca. 4mm) |
| 13. S | | |
| A | k. | SHAPE |
| 1 | | 1 = Ovate $2 = Oval$ $3 = Elliptical$ |
| B | i. ' | CHEEK |
| | | 1=Rounded 2=Angular |
| \mathbf{a}^{c} | :. l | BRUSH |
| <u></u> | | 1=Short 2=Medium 3=Long |
| 1 | | 1 = Not Collared 2 = Collared |
| _ |), (| CREASE |
| 1 | | 1 = Width 60% or less of Kernel 2 = Width 80% or less of Kernel 1 = Depth 20% or less of Kernel 2 = Depth 35% or less of Kernel |
| | | 3 = Width Nearly as Wide as Kernel 3 = Depth 50% or less of Kernel 3 = Depth 50% or less of Kernel |

| 13. | SEED: (continued) E. COLOR | |
|-----|--|--|
| 2 | $1 = \text{White} \qquad 2 = \text{Amber} \qquad 3 = \text{Red}$ | 4 = Other (specify) |
| | F. TEXTURE | |
| 1 | 1=Hard 2=Soft | |
| _ | G. PHENOL REACTION (see instructions): | |
| 0 | 1 = Ivory $2 = Fawn$ $3 = Light Brow$ | wn 4 = Dark Brown 5 = Black |
| | DISEASE: (0=Not Tested; 1=Susceptible; ASE INDICATE THE SPECIFIC RACE OR STRAIN TESTED | 2=Resistant; 3=Intermediate; 4=Tolerant) 5=moderately resistant 6=moderately susceptible |
| 2 | Stem Rust (Puccinia graminis f. sp. tritici) Field races | Leaf Rust (Puccinia recondita f. sp. tritici) Field races |
| 0 | Stripe Rust (Puccinia striiformis) | 1 Loose Smut (Ustilago tritici) |
| 3 | Tan Spot (Pyrenophora tritici-repentis) | Plag Smut (Urocystis agropyri) |
| 0 | Halo Spot (Selenophoma donacis) | Common Bunt (Tilletia tritici or T. laevis) |
| 0 | Septoria nodorum (Glume Blotch) | Dwarf Bunt (Tilletia controversa) |
| 0 | Septoria avenae (Speckled Leaf Disease) | (Karnal Bunt (Tilletia indica) |
| 5 | Septoria tritici (Speckled Leaf Blotch) Field races | Powdery Mildew (Erysiphe graminis f. sp. tritici) Field races |
| 0 | Scab (Fusarium spp.) | 0 Snow Molds |
| 0 | Black Point (Kernel Smudge) | Common Root Rot (Fusarium, Cochliobolus and Bipolaris spp. |
| 0 | Barley Yellow Dwarf Virus (BYDV) | Rhizoctonia Root Rot (Rhizoctonia solani) |
| 2 | Soilborne Mosaic Virus (SBMV) Field races | Black Chaff (Xanthomonas campestris pv. translucens) |
| 0 | Wheat Yellow (Spindle Streak) Mosaic Virus Field races | Bacterial Leaf Blight (Pseudomonas syringae pv. syringae) |
| 5 | Wheat Streak Mosaic Virus (WSMV) Field races | Other (specify) |
| 5 | Other (specify) Spindle streak mosaic virus (SSMV) | Other (specify) |
| | Other (specify) | Other (specify) |
| | Other (specify) | Other (specify) |

Exhibit D. Additional Description of Jagalene

Jagalene is a hard red winter wheat bred and developed by AgriPro Wheat. Jagalene is semidwarf in plant height and has white chaff at maturity. It has medium maturity and excellent straw strength. Jagalene is resistant to Stem Rust, Leaf Rust and Soil Borne Wheat Mosaic Virus. It is moderately resistant to Speckled leaf blotch, Wheat Spindle Streak Mosaic Virus and Wheat Streak Mosaic Virus. Jagalene has an intermediate reaction to Tan spot and is moderately susceptible to Powdery Mildew. Jagalene is susceptible to Hessian Fly and the wheat curl mite.

Juvenile growth habit is semierect. Plant color at boot stage is blue green. Auricle anthocyanin and auricle hairs are present. Flag leaf at boot stage is erect and twisted. Waxy bloom is present on the head, stem and flag leaf sheath. Anther color is yellow. Head shape is strap and awned. Glumes are glabrous, medium in width and long in length with oblique shoulders and acuminate beaks. Seed shape is ovate. Brush hairs are medium in size. Seed crease depth is shallow and width is narrow. Seed cheeks are rounded.

Jagalene is adapted to the Great Plains states of Colorado, South Dakota, Nebraska, Kansas, Oklahoma and Texas.

AGRIPRO

Plains Team Quality Summary

| | | | Color | 1010 | ~ | | , | 1 C | 1 tu | , m | ı m | , (m | 1.64 | · 10 | ı m | 4 | · m | . 67 | · m | ו רי | , (* | ı m | | r | 1 C | m | m | m | m | m | 4 | т | ю | m | . 60 | т | m | m | 60 |
|---------------------|----------------|-----------|----------|------------|------------|--------------|-----------|-----------|-----------|-----------|--------|-----------|-----------|-----------|--------|-----------|-----------|-----------|-----------|-----------|-----------|---------|--------|-----------|-----------|-----------|-----------|---------|-----------|-----------|-----------|--------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| | <u>-</u> | O III | Toy | <u> </u> | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | Č | | - | - - | ~ | | ι, | , , | 1 (7) | ייי | י ניי | , m | · m | 7 | 3 | | ω, | | : m | . 7 | 1 (**) | · m | | t. | וא נ | (L) | m | e. | c, | ιŋ | 7 | m | 6.1 | m | ω, | c | 4 | 3 | m |
| | ≧ | | Crain | 5 | ~ | | í. | 4 | 4 | 4 | 4 | 4 | ζ. | 4 | ĸ | 4 | 4 | 4 | E | 4 | . tu | 4 | | " | 4 | 50 | n | ۰۷۰ | e | 4 | 4 | 4 | 4 | 33 | 4 | n | c | Э | 4 |
| | Baking Quality | <u>ب</u> | | _ | ¥ | | 55 | 5 | . 8 | 55 | 35 | 51 | 55 | 15 | 오 | 5 | <u>o</u> | 0 | 9 | 5 | Q | gg. | | 0 | . N | 'n | ώ | رة ا | 0 | 0 | 0 | 0 | 5 | 5 | 0 | 0 | 40 | 5 | 0 |
| | Sakin | Loaf | | Vol | ວ | | 26 | 8 | . % | 6 | 103 | 1045 | 96 | 93 | 2 | 97 | 100 | 106 | 95 | 104 | 89 | 963 | | 16 | 805 | 85 | 25 | 119 | 94 | 93 | 8 | 95 | 100 | 91 | 95 | 95 | 95 | 89 | 94 |
| | | × | 9 | ; | <u>~</u> | | 75 | 2 | , Š | 00 | . 2/ | 0 | 0 | 75 | 55 | 75 | 00 | 0 | 99 | 5 | λ. | 22 | | 9 | Š | 5 | 5 | 5 | 0 | S | 'n | 50 | 5 | 0 | Ś | 0 | S | 'n | 23 |
| | | M | Time | | mim | | m | 5.(| ý | 3, | ٣ | 3.00 | 3. | 33 | 3.2 | 3.7 | 4 | 4.0 | 3.4 | 3.7 | 3.2 | 3.8 | | 3,0 | 4.75 | 4.7 | 3.2 | 3.7 | 3.0 | 2.7 | 3.7 | 2.7 | 2.7 | 4.0 | 3.2 | 3.5 | 4.2 | 3.2 | 3.5 |
| | | | | | ~ | (m) | _ | _ | _ | _ | | _ | _ | _ | | | _ | | _ | _ | _ | | | _ | _ | _ | _ | | | | | | | | | | | | |
| | | | Abs | | % | ENE | 62.0 | 62.0 | 9.09 | 62.(| 61.0 | 64.0 | 91.0 | 62.0 | 60.3 | 60.5 | 99 | 63.0 | 62.0 | 62.0 | 59.0 | 61.8 | ER | 64.0 | 62.0 | 61.0 | 62.0 | 63.0 | 64.5 | 62.0 | 62.0 | 62.0 | 90.0 | 62.5 | 63.0 | 63.0 | 62.0 | 61.0 | 62.3 |
| | | | _: | | ~ | IAGAI | 42 | 965 | 20 | 45 | 61 | 86 | 13 | 34 | 32 | 33 | 요 | 6/ | ₽ | 77 | 35 | 33 | JAGGER | 33 | 9, | 200 | i.c. | ∞ | 2 | = | 4 | 6 | 6 | ∞ | Ģ | 7 | ∞ | 2 | 4 . |
| | ram | | Tol. | | HH | JA | 1042 | | 1150 | | • | 798 | | - | _ | | | | | 1022 | 1035 | 983 | J. | 78 | 926 | 103 | ≅ | 75 | 55 | 63 | 66 | 2 | 19 | 99 | 84 | 78 | 110 | 103 | ∞ |
| | Mixogram | Peak | | HI | Ż. | | 5.0 | 5.3 | 4.8 | 5.5 | 5.3 | 5.0 | 5.0 | 4.5 | 4.8 | 5.0 | 5.0 | 5.0 | 5.0 | 4.8 | 4.8 | 5.0 | | 5.3 | 5.5 | 5.3 | 5.5 | 5.8 | 5.3 | 5.3 | 5.0 | 5.3 | 5.5 | 5.0 | 5.3 | 5.3 | 5.0 | 5.3 | 5,3 |
| | _ | Peak Peak | | Time | ii ii | | 3.75 | 5.00 | 5.75 | 3.50 | 3.75 | 3.00 | 3.00 | 3.75 | 3.25 | 3.75 | 4.25 | 4.00 | 3.50 | 4.00 | 3.25 | 3.83 | | 3.00 | 4.75 | 4.75 | 3.25 | 3.50 | 3.00 | 2.75 | 3.75 | 2.75 | 2.75 | 4.00 | 3.25 | 3.25 | 4.25 | 3.25 | 3.48 |
| ality | | | Ash | | | | | | | 0.530 | 0.450 | 0.476 | 0.468 | 0.442 | 0.492 | 0.474 | 0.468 | 0.442 | 0.436 | 0.450 | 0.434 | 0,464 | | | | | 0.564 | 0.494 | 0.456 | 0.478 | 0.388 | 0.472 | 0.484 | 0.380 | 0.400 | 0.420 | 0.450 | 0.454 | 0.453 |
| Saf Or | , | | | ı | ~ | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Flour/Wheat Onality | | Flr | | ΝId | % | | 73.0 | 71.7 | 70.8 | 69.5 | 67.3 | 71.7 | 71.1 | 9.69 | 71.9 | 70.6 | 69.3 | 70.5 | 70.7 | 70.3 | 70.1 | 70.5 | | 70.7 | 69.3 | 68.3 | 67.3 | 65.3 | 72.4 | 70.8 | 67.8 | 71.8 | 68.2 | 68.7 | 70.8 | 70.8 | 69.7 | 6.69 | 69.5 |
| Floir | | Norris | | Hard | | | 84 | 73 | 79 | 92 | 80 | 81 | 77 | 70 | 7.5 | 74 | 79 | 9/ | 78 | 72 | 99 | 11 | | 93 | 59 | 73 | 90 | 80 | 72 | 80 | 59 | 81 | 7.1 | 69 | 71 | 75 | 64 | 65 | 73 |
| | | | | | × | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | Flr | Prot | | 4%mb | | 10.7 | 11.0 | 9.6 | 12.3 | 12.8 | 13.4 | 12.8 | 11.3 | 11.2 | 11.2 | 12.7 | 12.8 | 12.5 | 12.7 | 10.8 | 11.9 | | 11.4 | 11.7 | 10.0 | 11.7 | 14.8 | 14.0 | 13.1 | 11.3 | 11.9 | 11.6 | 11.9 | 12.8 | 13.3 | 13.2 | 12.0 | 12.3 |
| | | Wht | | Prot | 14%mb 14%m | | 12.0 | 12.3 | 11.1 | 13.6 | 14.2 | 14.3 | 13.8 | 12.5 | 12.3 | 12.3 | 13.9 | 13.8 | 13.5 | 13.7 | 11.9 | 13.0 | | 13.0 | 13.1 | 11.6 | 13.3 | 15.8 | 14.9 | 14.2 | 12.6 | 12.7 | 12.7 | 12.9 | 13.5 | 14.1 | 14.4 | 12.9 | 13.4 |
| | | | Year-Loc | — ; | 14 | | 1998 - SK | 1999 - QK | 2000 - EB | 2000 - VT | 2000 - | 2001 - DL | 7001 - DT | 2001 - EB | 2001 - | 2001 - HP | 2001 - JP | 2001 - PD | 2001 - PD | 2001 - VD | 2001 - VD | Average | | 1998 - SK | 1999 - QK | 2000 - EB | 2000 - VT | 2000 - | 2001 - DL | 2001 - DL | 2001 - EB | 2001 - | 2001 - HP | 2001 - JP | 2001 - PD | 2001 - PD | 2001 - VD | 2001 - VD | Average |

Ratings: 1-2=Excellent 3-4=Good 5-6=Acceptable 7-8=Questionable 9=Unacceptable

200200160

Three Year Summary for Yield (bu/a) of selected varieties tested in the AgriPro commercial hard red winter wheat testing program

| <u>Variety</u> | | <u>EAST</u> | | | | <u>West</u> | | |
|----------------|-------------|-------------|-------------|----------------|-------------|-------------|-------------|----------------|
| | <u>1999</u> | <u>2000</u> | <u>2001</u> | <u>Average</u> | <u>1999</u> | 2000 | <u>2001</u> | <u>Average</u> |
| Jagger | 60 | 58 | 60 | 59 | 68 | 46 | 65 | 60 |
| 2137 | 52 | 52 | 53 | 52 | 65 | 44 | 54 | 54 |
| Jagalene | 59 | 65 | 56 | 60 | 69 | 51 | 69 | 63 |
| Cutter | 54 | 61 | 60 | 58 | 70 | 44 | 65 | 60 |
| Dumas | 55 | 63 | 51 | 56 | 64 | 47 | 58 | 56 |
| Coronado | 56 | 53 | 53 | 54 | 63 | 45 | 57 | 55 |
| Hondo | 48 | 49 | 52 | 50 | * | * | * | * |
| Big Dawg | 61 | 50 | 57 | 56 | * | * | * | * |
| Thunderbolt | * | * | * | * | 63 | 42 | 56 | 54 |
| Ogallala | * | * | * | * | 63 | 42 | . 58 | 54 |

East, 1999 = Salina

East, 2000 = Salina, Nardin, Partridge

East, 2001 = Salina, Junction City, Everest, Partridge, Winfield

West, 1999 = Paxton, Quinter, Hugoton, Otis

West, 2000 = Paxton, Quinter, Hugoton, Goodland, Otis

West, 2001 = Paxton, Otis, Goodland, Manter, Hugoton

AgriPro Wheat comparison chart for agronomic, disease and Insect reactions for currently grown hard winter wheets adapted to the Southern Plains. Rating range from 1 (excellent) to 9 (worst)

| <u>¥</u> ≈ | B ∵ 2 | ထေးတက | ဘထတ ။ | သောထာဏ | 5 60 | တဆေးဆ |) 00 | . 00 00 | · (| 7 6 | 6 h | מיז - | α (| ו תב |
|---------------------------|------------------------|----------------------------------|-------------------------------|------------------------|----------------|--------------------|-------------------|-----------------|-------------------|-------------|-------------------|--------------|--------------|----------|
| PM 5 | a 15 a | ~ m m · | o 62 4 € | v ~ r | - 5 0 | ភ | 2 | W | | 4 C/ | ~ σ | ော | ധ | N 0 |
| lan spot 5 | υ τo 4 ⋅ | ፋጥጭኖ |) w w < | * OC C | · 100 | ω Φ 4 | 4 | æ 4 | ر م | വാ | 5. ~ | د د | ~ r | ~ a |
| S. trit | ÷ ω ຕ • | - 60 60 | 11-40 | ကေးထ | ~ | N 4 4 | ಣ | ~ ⊕ | 40 | o 4 | 7 | wι | Þ١ | - 1- |
| WSMV 8 | o Co 🚓 Li | 9 4 . B . L | 0 4 4 | . 4- r∪ | ល | o 4 4 | 4 | 40 | 4 1 | | മ | ≻ 0 | o w | > т; |
| SSMV 4 | · * * * | ነ ሰ 4 ሲ | மம− | លស | សេះ | ာကကေး | 4 | 9 ~ | 4.6 | 4 | ာထ | 6 0 0 | n da | . 4 |
| SBMV 1 | . , , , , , | | ಌ ಱ ← | ∞ | , . | - ← ຄ | - | 80 / - | ٠. | ₩ (| ာထ | oc w | , c | N |
| Leaf R 4 3 | 9 \$ 1 | ಬಣ⊢ | ខេត្ត | 4 ۲. | ∞ 1~ | . | 8 | සා ස | ۰ 9 | יט כו | b 00 | 00 LC | , 6 0 | ঘ |
| Stem R 5 3 | ೧೮4 | 8 th ∕ | ত ধ ধ | ব ব | - বিংগ |) ro ro | TO I | ω 4 : | ~ 4 | φ < | . rv | 4. 10 | * | en |
| Protein 4 5 | æ ፋ ro | 446 | 4 ት r c | ഥന [.] | 4 4 | . 4 ю | က၊ | ~ LS I | v 4 | 40 t | - (C) | ம ம | 7 | s, |
| Al. Tol 5 8 | ro 4 '0 | ಹದಾದ | သေးထာက | r- r 1 | c 4 | 44 | 4 (| ာထ ေ | n co | ကေးထ | , es 1 | ထထ | æ | € |
| WH 6 5 | | დ ი ი ი · | 4 4 W | ကက | ကယ | ខេត | တဂ | o Co L | D 4 | æ 4 | 4 - | 4 W | ဗ | 2 |
| Shatter 4 4 | a 4 r | ကတက | იდ 4 : | ဝတဗေ | o 4 | ເລ ແລ | 9 5 | t 60 ⊀ | t ro | ro 4 | 4 * | t 4 | 4 | 4 |
| Col. Leng. 5 · | വവാ | ი 4 თ • | 1 ro 4 j | 2(a) r | 2 2 | ជាជា | 4 - | ן עס ע | . a. | დ 4 | ro e | ខ្មា | 4 | ഗ |
| | ର ୧୯୯ । | থ ব ব ব | ა დ. 4. მ | യെ | 0 0 | 6/ TO | 4 4 | . 10 A | - co - | v 4 | ro r | 9 | 4 | 4 |
| rel. Mat. 3 5 | າຕຕາ | 4 4 4 4 | ւտտո | വവാധ |) t~ | ညယ | 20 | 14 rc | · 55 · | 4 13 | w ie | î - | ლ. | 4 |
| ιo | מממ | থ বং ব | - 01 10 2 | 122 | 2 | 2 23 | د 4 | . m w | · 03 • | * + | សព | ۸ د | -d : | ro Co |
| | ነዋድ | oro ⊠. 4 | - ପ ୩ ର | 100 | 2 | 01 m | സന | 60 4 | е • | 4 V | ស ស | ı Çı | ro c | 'n |
| HICKOCK Dumas PECOS | CORONADO ORO BLANCO | TOMAHAWK THUDERBOLT LAREDO | OGALLALA HONDO LONGHORN | THUNDERBIRD ABILENE | BIG DAWG | Jagalene Cufter | JAGGER TAM 107 | CUSTER 2137 | DOMINATOR 2474 | TAM 110 | AKRON ALLIANCE | ARAPAHOE | Fraine Ked | OBale |

Ratings taken by John Moffatt, David Worrall and Rollin Sears, AgriPro Wheat winter wheat breeders



Hard Winter Wheat PROfile

AgriPro Variety 'Jagalene'

Abijene/Jagger

Excellent "all purpose" Wheat




| Variety Features | |
|------------------|-----------|
| Plant Height | semidwari |
| Emergence | Average |
| Fall Cover | Average |
| Winter Survival | Very Good |
| Spring Cover | Medium |
| Coleoptile | Viedium |
| Straw Strength | Excellent |
| Maturity | Medium |

| Head and G | Grain |
|-------------------|-----------|
| Туре | Awned |
| Seed Size | Large |
| Test Weight | Very Good |
| Drydown | Quick |
| Shatter Tolerance | Good |
| Chaff Color | White |
| Quality | Excellent |

Production Potential ★ Widely adapted ★ Excellent drought tolerance ★ Great disease package

- 보 Large kernels & top test weights
- k Excellent milling & baking quality



| Greenbug | 745,450,7755 | *1.402.4623724255 | | 持续的 177 | | |
|---------------------|--------------|--|---------------------|--|--|-------------------|
| Soil Borne Complex | E E E | Notes of the second | Sacrati es para | | All participations | 153 |
| Barley Yellow Dwarf | | | | | | |
| Wheat Streak Mosaic | 254599 | ###################################### | The Ry of The State | POST STOCKEN | | |
| Powdery Mildew | 11.7FT (6) | restriction of | | | | |
| Tan Spot | 4946 | | | | | Allerad Straus |
| Leaf Blotch | 100000 | 5. 存取的 3. 3. 467. 1 | ann an things | - 12 CONTRACT PARTY - 12 C | | |
| Stem Rust | (d) (d) | | #100 Politicals | | Conservation of the con- | [漢] |
| Leaf Rust | 2302 B | e segunda alame | | ALL STATES | e de la companya de l | iasse Lore |
| | 'n | 2 | .4 | 5 | 9 | 10 |

Susceptible

| 10,01) Agril | ⁵ro Yield Summai | ry |
|--------------|-----------------------------------|----------------|
| Central Plai | ns (25 locations) | |
| Yield | TW | |
| 63 | 60 | |
| 60 | 59 | |
| 54 | 58 | |
| | | |
| | | |
| | Central Plai Yield 63 60 | 63 60 60 59 |

© 2001 AgriPro Wheet. AgriPro Wheet is a business unit of Advanta, USA. All railings, including plant height and seed size, may vary with growing conditions.

Intermediate

Exhibit E. Statement of the Basis of Applicant's Ownership

The variety for which Plant Variety Protection is hereby sought was developed by Dr. John Moffatt, an employee of AgriPro Wheat, a business unit of Advanta USA. By agreement between employees and AgriPro Wheat, all rights to any invention, discovery, or development made by the employee while employed by AgriPro Wheat, were assigned to AgriPro Wheat with no rights of any kind pertaining to 'Jagalene' being retained by the employees.

By contractual agreement the variety 'Jagalene' was purchased from AgriPro Wheat, a business unit of Advanta USA in June of 1996 and is currently owned by Monsanto Company.